

AN EDGE-COLOURED VERSION OF DIRAC'S THEOREM

ALLAN LO

*School of Mathematics, University of Birmingham,
Birmingham, B15 2TT, UK*

ABSTRACT. Let G be an edge-coloured graph. Suppose that for every vertex v there are at least $\delta^c(G)$ distinct colours on edges incident to v . We say that G is properly coloured if no two adjacent edges have the same colour. In this paper, we show that every edge-coloured graph G with $\delta^c(G) \geq 2|G|/3$ contains a properly coloured 2-factor. Furthermore, we show that for any $\varepsilon > 0$ there exists an integer n_0 such that every edge-coloured graph G with $|G| = n \geq n_0$ and $\delta^c(G) \geq (2/3 + \varepsilon)n$ contains a properly coloured cycle of length ℓ for every $3 \leq \ell \leq n$. This result is best possible in the sense that the statement is false for $\delta^c(G) = (2/3 - \varepsilon)n$.

1. INTRODUCTION

An *edge-colouring* c of a graph G is an assignment of colours to the edges of G . An *edge-coloured graph* is a graph G with an edge colouring c of G . An edge-coloured graph G is *properly coloured* if no two adjacent edges of G have the same colour. If all edges have distinct colours, then G is *rainbow*. If all edges have the same colour, then G is *monochromatic*.

Grossman and Häggkvist [8] were the first to give a sufficient condition for the existence of properly coloured cycles in edge-coloured graphs with two colours. Later on, Yeo [14] extended the result to edge-coloured graphs with any number of colours. A natural question is to ask what guarantees the existence of properly coloured Hamiltonian paths and cycles. For an edge-coloured complete graph K_n^c on n vertices, Beng-Jensen, Gutin and Yeo [4] proved that if K_n^c contains a properly coloured 2-factor, then K_n^c also contains a properly coloured Hamiltonian path. Later, Feng, Giesen, Guo, Gutin, Jensen and Rafiey [7] improved this result and gave a necessary and sufficient condition for the existence of a properly coloured Hamiltonian path. A graph G is said to be a *1-path-cycle* if G is a vertex-disjoint union of at most one path and a number of cycles. Feng et al. [7] showed that K_n^c contains a spanning properly coloured 1-path-cycle if and only if K_n^c contains a properly coloured Hamiltonian path.

E-mail address: s.a.lo@bham.ac.uk.

Date: January 1, 2013.

Key words and phrases. proper edge-coloring, 2-factor, Hamiltonian cycle.

The author was supported by the ERC, grant no. 258345.

Given an edge-coloured complete graph K_n^c on n vertices, $\Delta_{\text{mon}}(K_n^c)$ is defined to be the maximum number of edges of the same colour incident to the same vertex. In other words, $\Delta_{\text{mon}}(K_n^c) = \max \Delta(H)$ over all monochromatic subgraphs $H \subseteq K_n^c$. Bollobás and Erdős [5] proved that if $\Delta_{\text{mon}}(K_n^c) \leq n/69$, then K_n^c contains a properly coloured Hamiltonian cycle. They further conjectured that $\Delta_{\text{mon}}(K_n^c) < \lfloor n/2 \rfloor$ suffices. Their result was subsequently improved by Chen and Daykin [6], Shearer [13] and Alon and Gutin [1]. In [11], the author showed that for any $\varepsilon > 0$ every K_n^c with $\Delta_{\text{mon}}(K_n^c) < (1/2 - \varepsilon)n$ contains a properly coloured Hamiltonian cycle, provided n is large enough. Therefore, the conjecture of Bollobás and Erdős is true asymptotically. For a survey regarding properly coloured subgraphs in edge-coloured graphs, we recommend Chapter 16 of [3].

Given an edge-coloured graph G (not necessarily complete), the *colour degree* $d^c(v)$ of a vertex v is the number of distinct colours of edges incident to v . The *minimum colour degree* $\delta^c(G)$ of an edge-coloured graph G is the minimum $d^c(v)$ over all vertices v in G . Li and Wang [9] proved that G contains a properly coloured path of length $2\delta^c(G)$ or a properly coloured cycle of length at least $2\delta^c(G)/3$. In [10], the author improved this result by showing that G contains a properly coloured path of length $2\delta^c(G)$ or a properly coloured cycle of length at least $\delta^c(G) + 1$. In the same paper, the author showed that every connected edge-coloured graph G contains a properly coloured Hamiltonian cycle or a properly coloured path of length $6\delta^c(G)/5 - 1$. Furthermore, the author also conjectured the following.

Conjecture 1.1 ([10]). *Every connected edge-coloured graph G contains a properly coloured Hamiltonian cycle or a properly coloured path of length $\lfloor 3\delta^c(G)/2 \rfloor$.*

If the conjecture is true, then every edge-coloured graph G with $\delta^c(G) \geq 2|G|/3$ contains a properly coloured Hamiltonian cycle. In this paper, we prove the following weaker result that $\delta^c(G) \geq 2|G|/3$ implies a properly coloured 2-factor.

Theorem 1.2. *Every edge-coloured graph G with $\delta^c(G) \geq 2|G|/3$ contains a properly coloured 2-factor.*

We also show that if $\delta^c(G) \geq (2/3 + \varepsilon)|G|$, then G does indeed contain a properly coloured Hamiltonian cycle. Moreover, G is properly coloured pancyclic.

Theorem 1.3. *For any $\varepsilon > 0$, there exists an integer n_0 such that every edge-coloured graph G with $\delta^c(G) \geq (2/3 + \varepsilon)|G|$ and $|G| \geq n_0$ contains a properly coloured cycle of length ℓ for all $3 \leq \ell \leq |G|$.*

Given a subgraph H in G , we write $G - H$ for the (edge-coloured) subgraph obtained from G by deleting all edges in H . Given an edge-coloured graph G , let $\delta_1^c(G)$ be the minimum $\delta(G - H)$ over all monochromatic subgraphs H in G . Note that $\delta_1^c(G) \geq \delta^c(G) - 1$. Also for an edge-coloured complete graph K_n^c , $\delta_1^c(K_n^c) = n - 1 - \Delta_{\text{mon}}(K_n^c)$. We prove the following stronger statement, which implies Theorem 1.3.

Theorem 1.4. *For any $\varepsilon > 0$, there exists an integer n_0 such that every edge-coloured graph G with $\delta_1^c(G) \geq (2/3 + \varepsilon)|G|$ and $|G| \geq n_0$ contains a properly coloured cycle of length ℓ for all $3 \leq \ell \leq |G|$.*

We now outline the proof of Theorem 1.4 for the case when $\ell = |G|$ i.e. the existence of a properly coloured Hamiltonian cycle. The proof is divided into two main steps. In the first step, we find (by Lemma 4.1) a small ‘absorbing cycle’ C in G such that given any small number of vertex-disjoint properly coloured paths $P_1, P_2, \dots, P_k$ in G with $V(C) \cap V(P_i) = \emptyset$ for each $i \leq k$, there exists a properly coloured cycle C' in G with $V(C') = V(C) \cup \bigcup_{1 \leq i \leq k} V(P_i)$. This step can be viewed as a properly edge-coloured version of the absorption technique introduced by Rödl, Ruciński and Szeemerédi [12]. Since the original absorption technique did not consider edge-coloured graphs, several new ideas are needed for this generalization. We believe that there is further potential for this adaptation of the absorption technique. For instance, a similar argument was also used in [11]. In the second step, we remove the vertices of C from G and let G' be the resulting graph. Since C is small, we may assume that $\delta_1^c(G') \geq (2/3 + \varepsilon')|G'|$ for some small $\varepsilon' > 0$. Then we find a properly coloured 2-factor in G' using Lemma 3.3 such that every cycle has length at least $\varepsilon'|G'|/2$. Hence, $V(G')$ can be covered by at most $2\varepsilon'^{-1}$ vertex-disjoint properly coloured paths $P_1, P_2, \dots, P_k$. By the ‘absorbing’ property of C , there is a properly coloured cycle C' with $V(C') = V(C) \cup \bigcup_{1 \leq i \leq k} V(P_i) = V(G)$. Therefore, C' is a properly coloured Hamiltonian cycle as required.

The paper is organised as follows. In the next section, we set up some basic notation and give the extremal example for Conjecture 1.1. Section 3 will be devoted to finding properly coloured 2-factors, where we prove Theorem 1.2. The absorbing cycle is constructed in Section 4. Finally, we prove Theorem 1.4 in Section 5.

2. NOTATION AND THE EXTREMAL EXAMPLE

Throughout this paper unless stated otherwise, G is assumed to be an edge-coloured graph with edge-colouring c . Hence, $c(xy)$ is the colour of the edge xy . For $v \in V(G)$, we denote by $N_G(v)$ the neighbourhood of v in G . If the graph G is clear from the context, we omit the subscript.

Given a vertex set $U \subseteq V(G)$, write $G[U]$ to be the (edge-coloured) subgraph of G induced by U . We write $G \setminus U$ for the graph obtained from G by deleting all vertices in U . For edge-disjoint graphs G and H , we denote by $G + H$ the union of G and H . We write $G - H + H'$ to mean $(G - H) + H'$.

Each path P is assumed to be directed. Hence, the paths $v_1v_2 \dots v_\ell$ and $v_\ell v_{\ell-1} \dots v_1$ are considered to be different. Given a path $P = v_1v_2 \dots v_\ell$ and a vertex $x \in N(v_1) \setminus V(P)$, we denote by xP the path $xv_1v_2 \dots v_\ell$. Similarly, given vertex-disjoint paths $P_1, \dots, P_s$, then we define $P_1 \dots P_s$ to be the concatenation of $P_1, \dots, P_s$.

Let H be a union of vertex-disjoint directed cycles. For each $y \in V(H)$, let C_y be the cycle in H that contains y . Denote by y_+ and y_- the successor and ancestor of y in C_y respectively. Write y_- and y_{++} for $(y_-)_-$ and $(y_+)_+$. For distinct vertices y and z in a cycle C in H , define yC^+z and

yC^-z to be the paths $yy_+ \dots z_-z$ and $yy_- \dots z_+z$ in C respectively. If $y = z$, then set $yC^+y = y = yC^-y$.

In [10], the author gave a construction to show that Conjecture 1.1 is best possible for infinitely many values of $|G|$ and $\delta^c(G)$. The same construction also shows that Theorem 1.2 is best possible. We include it here for completeness.

Proposition 2.1. *For each n , there exists a connected edge-coloured graph G on n vertices with $\delta^c(G) < 2|G|/3$ which do not contain vertex-disjoint properly coloured cycles spanning greater than $3\delta^c(G)/2$ vertices. In particular, G does not contain a properly coloured 2-factor or a properly coloured Hamiltonian cycle. Moreover, all properly coloured paths in G have length at most $3\delta^c(G)/2$.*

Proof. Let n and δ be integers with $3\delta < 2n$. Let $X = \{x_1, x_2, \dots, x_\delta\}$ and Y be vertex sets with $|Y| = n - \delta$ and $X \cap Y = \emptyset$. Define G to be the edge-coloured graph on the vertex set $X \cup Y$ as follows. Let $G[X]$ be a rainbow complete graph and let $G[Y]$ be a set of independent vertices. For each $1 \leq i \leq \delta$, add an edge of new colour c_i between x_i and y for each $y \in Y$. By our construction, $d^c(v) = |X| = \delta$ for all $v \in V(G)$ and so $\delta^c(G) = \delta$.

Let C be a properly coloured cycle in G . Orient C into a directed cycle. Note that every vertex $y \in V(C) \cap Y$ must be followed by two consecutive vertices in X , so $|X \cap V(C)| \geq 2|Y \cap V(C)|$. Therefore, if $\mathcal{C}$ is a collection of vertex-disjoint properly coloured cycles in G , then $\mathcal{C}$ spans at most $|X| + |X|/2 = 3\delta/2$ vertices. The ‘moreover’ statement is proved by a similar argument. $\square$

3. PROPERLY COLOURED 2-FACTORS

First we prove Theorem 1.2; that is, every edge-coloured graph with $\delta^c(G) \geq 2|G|/3$ contains a properly coloured 2-factor. Here we present a sketch of the proof. Suppose that G is an edge-coloured graph on n vertices. Let H be a properly coloured subgraph in G consisting of vertex-disjoint cycles such that $|H|$ is maximal. If $V(H) = V(G)$, then we are done. Hence we may assume that there exists $x \in V(G) \setminus V(H)$. Define the *colour neighbourhood* $N^c(x)$ of x in G to be a maximal subset of neighbours of x in G such that $c(xy) \neq c(xz)$ for every distinct $y, z \in N^c(x)$. (The choice on $N^c(x)$ will be specified later.) Assume that we are in the ideal case that $N^c(x) \cap V(H) = \emptyset$. If $c(xy) \neq c(yz) \neq c(xz)$ for distinct $y, z \in N^c(x)$, then $xyzx$ is a properly coloured triangle and so $H + xyzx$ contradicts the maximality of $|H|$. Hence, we may assume that for every $y, z \in N^c(x)$, either yx is not an edge, $c(zx) = c(xy)$ or $c(zx) = c(xz)$. By an averaging argument, there exists a vertex $y_0 \in N^c(x)$ such that the number of $z \in N^c(x) \setminus y_0$ such that either $y_0z \notin E(G)$ or $c(y_0z) = c(xy_0)$ is at least $(|N^c(x)| - 1)/2 \geq (\delta^c(G) - 1)/2$. Recall that there are at least $\delta^c(G) - 1$ neighbours v of y_0 with $c(y_0v) \neq c(xy_0)$. This means that $n = |V(G)| \geq (\delta^c(G) - 1)/2 + (\delta^c(G) - 1) + |\{x, y_0\}| > 3\delta^c(G)/2$ and so $\delta^c(G) < 2n/3$.

Proof of Theorem 1.2. Let G be an edge-coloured graph on n vertices with edge-colouring c . Set $\delta = \delta^c(G)$ and $V = V(G)$. Suppose that G does not

contain a properly coloured 2-factor. We will show that $\delta < 2n/3$. This is trivial if $n \leq 2$, so we may assume that $n \geq 3$. By deleting edges in G , we may assume that G is edge-minimal such that any additional edge deletion would lead to a decrease in $d^c(v)$ for some vertex v in G . If a monochromatic subgraph G' in G is not isomorphic to a disjoint union of stars, then there exists an edge uv in G' with $d_{G'}(u), d_{G'}(v) \geq 2$. Let $G'' = G - uv$ be the edge-coloured graph obtained from G after removing the edge uv . Note that $d_{G''}^c(x) = d_G^c(x)$ for all vertices $x \in V(G)$. This contradicts the edge-minimality of G . Therefore, every monochromatic subgraph G' in G is a disjoint union of stars. Let H be a properly coloured subgraph in G consisting of vertex-disjoint cycles such that $|H|$ is maximal. Note that $|H| < n$. Arbitrarily orient each cycle in H into a directed cycle.

Fix $x \in V \setminus V(H)$. Let $N^c(x)$ be a maximal subset of neighbours of x in G such that $c(xy_1) \neq c(xy_2)$ for every distinct $y_1, y_2 \in N^c(x)$. Note that $|N^c(x)| = d^c(x)$. If there are at least two vertices y such that the edge xy has the same colour, then we choose y to be in $N^c(x)$ satisfying the following order of preferences (if there are still at least two choices for y , pick one arbitrarily):

- (a) $y \notin V(H)$;
- (b) $y \in V(H)$ such that $c(xy) = c(yy_+)$;
- (c) $y \in V(H)$ such that $c(xy) = c(yy_-)$;
- (d) $y \in V(H)$ such that $xy_+ \notin E(G)$;
- (e) $y \in V(H)$ such that $c(xy) \neq c(xy_+)$;
- (f) $y \in V(H)$ such that $c(xy) = c(xy_+)$.

Note that if $y \in N^c(x)$ satisfies property (f), then $c(xy') = c(xy)$ for every vertex y' in C_y . Define $N_{(a)}^c(x)$ to be the set of vertices in $N^c(x)$ chosen due to (a), and define $N_{(b)}^c(x), \dots, N_{(f)}^c(x)$ similarly. Next, set

$$\begin{aligned} W &= N_{(a)}^c(x), \\ R' &= \{y_+ : y \in N_{(b)}^c(x) \cup N_{(d)}^c(x) \cup N_{(e)}^c(x) \cup N_{(f)}^c(x)\}, \\ S' &= \{y_- : y \in N_{(c)}^c(x)\}, \\ R &= R' \setminus S', \quad S = S' \setminus R', \quad T = R' \cap S'. \end{aligned}$$

Note that

$$|R| + |S| + 2|T| + |W| = |N^c(x)| \geq \delta. \quad (3.1)$$

If $y \in R' \cup S'$, then $y \in V(H)$. Since H consists of properly coloured cycles, we deduce that

- (i) if $y \in R'$, then $y_- \in N^c(x)$ and $c(y_-y_{--}) \neq c(xy_-)$, and
- (ii) if $y \in S'$, then $y_+ \in N^c(x)$ and $c(y_+y_{++}) \neq c(y_+y) = c(xy_+)$.

Note that (ii) is implied by the definitions of S' and $N_{(c)}^c(x)$. If $y \in R'$, then $y_- \in N_{(b)}^c(x) \cup N_{(d)}^c(x) \cup N_{(e)}^c(x) \cup N_{(f)}^c(x) \subseteq N^c(x)$. If $y_- \in N_{(b)}^c(x)$, then $c(y_-y_{--}) \neq c(y_-y) = c(xy_-)$ since C_{y_-} is a properly coloured cycle. If $y_- \in N_{(d)}^c(x) \cup N_{(e)}^c(x) \cup N_{(f)}^c(x)$, then $c(xy_-) \neq c(y_-y_{--})$ or else y_- should be in $y_- \in N_{(c)}^c(x)$. Hence, (i) is true.

Let F be the edge-coloured subgraph of G induced by the vertex set $W \cup R \cup S \cup T$. For $y \in V(F)$, define the vertex colour $c(y)$ to be

$$c(y) = \begin{cases} c(xy) & \text{if } y \in W, \\ c(yy_+) & \text{if } y \in R, \\ c(yy_-) & \text{if } y \in S, \\ c_0 & \text{if } y \in T, \end{cases}$$

where c_0 is a new colour that does not appear in G . In the following claim, we investigate the colours of the edges of F .

Claim 3.1. *If $yz \in E(F)$, then $c(yz) = c(y)$ or $c(yz) = c(z)$. In particular, $G[T]$ is empty.*

Proof of claim. Suppose the claim is false, so there exists an edge $yz \in E(F)$ with $c(yz) \neq c(y)$ and $c(yz) \neq c(z)$. We will show that there is a properly coloured subgraph H' in G consisting of vertex-disjoint cycles with $V(H') = V(H) \cup \{x, y, z\}$ contradicting the maximality of $|H|$.

If $y, z \in W \subseteq N^c(x)$, then $c(xy) \neq c(xz)$. Since $c(xy) = c(y) \neq c(yz) \neq c(z) = c(xz)$, then $xyzx$ is a properly edge-coloured triangle in G . Hence $H' = H + xyzx$ contradicts the maximality of $|H|$.

If $y \in W$ and $z \in R$, then $c(xy) = c(y) \neq c(yz) \neq c(z) = c(zz_+)$. By (i) we have $z_- \in N^c(x)$ and $c(xz_-) \neq c(z_-z_{--})$. Moreover, $c(xz_-) \neq c(xy)$ as $y, z_- \in N^c(x)$. Therefore, $C' = xyzC_z^+z_-x$ is a properly coloured cycle. We obtain a contradiction by setting $H' = H - C_z + C'$. Similarly, if $y \in W$ and $z \in S$, then we obtain a contradiction by considering $C'' = xyzC_z^-z_+x$ instead of C' . If $y \in W$ and $z \in T$, then $c(xy) = c(y) \neq c(yz)$. Since C_z is a properly coloured cycle, $c(yz) \neq c(zz_+)$ or $c(yz) \neq c(zz_-)$. Recall that $T = R' \cap S'$, so $c(xz_-) \neq c(z_-z_{--})$ and $c(xz_+) \neq c(z_+z_{++})$ by (i) and (ii). Also, $y, z_-, z_+ \in N^c(x)$, so $c(xz_-) \neq c(xy) \neq c(xz_+)$. Hence, $H - C_z + C'$ or $H - C_z + C''$ would imply a contradiction.

Therefore, we may assume that $y, z \in R' \cup S' \subseteq V(H)$. In order to prove the claim, it is enough to show that there exist at most two vertex-disjoint properly coloured cycles C', C'' spanning $\{x\} \cup V(C_y) \cup V(C_z)$ (which would then imply that $H' = (H - C_y - C_z) + C' + C''$ is a union of properly coloured cycles with $|H'| = |H| + 1$, a contradiction).

Suppose that C_y and C_z are distinct. If $y, z \in R'$, then by (i) we have

$$y_-, z_- \in N^c(x), \quad c(y_-y_{--}) \neq c(xy_-), \quad c(xz_-) \neq c(z_-z_{--}). \quad (3.2)$$

Since $y_-, z_- \in N^c(x)$, we have $c(xy_-) \neq c(xz_-)$. Note that $c(y) \neq c(yz) \neq c(z)$, so

$$c(yy_+) \neq c(yz) \neq c(zz_+) \quad (3.3)$$

Hence, $C' = xy_-C_y^-yzC_z^+z_-x$ (see Figure 1 (A)) is a properly coloured cycle with vertex set $V(C_y + C_z) \cup \{x\}$. Notice that C' is a properly coloured cycle if both (3.2) and (3.3) hold. If $y, z \in R'$, then (3.2) holds by (i). Therefore, we may assume that without loss of generality that $y \in S$ or $y \in T$ with $c(yy_+) = c(yz)$. Since $T \subset S'$ and the cycle C_y is properly coloured, we have $y \in S'$ and $c(yy_-) \neq c(yy_+) = c(yz)$. By (ii) and reversing the orientation on C_y , we have $y_- \in N^c(x)$, $c(y_-y_{--}) \neq c(xy_-)$ and $c(yy_+) \neq c(yz)$. Similarly,




FIGURE 1. Properly coloured cycles used in the proof of Claim 3.1

by reversing the orientation on C_z if necessary, we may assume that both (3.2) and (3.3) hold and so we derive a contradiction.

Finally, suppose that $C = C_y = C_z$. If $y, z \in R$, then $c(yy_+) = c(y) \neq c(yz) \neq c(z) = c(zz_+)$. In particular, $y_+ \neq z$ and $z_+ \neq y$. By (i), $xy_-C^-zyC^+z_-x$ (see Figure 1 (B)) is a properly coloured cycle with vertex set $V(C) \cup \{x\}$, so we are done. If $y \in R$ and $z \in S$, then $y_- \neq z$, otherwise xyC^+z is a properly coloured cycle by (i) and (ii). However, both $xy_-C^-z_+x$ and yC^+zy (see Figure 1 (C)) are properly coloured cycles spanning $\{x\} \cup V(C)$. If $y, z \in S$, then by reversing the orientation of the cycle C we conclude that $y, z \in R$ and so we are done. If $y \in T$ or $z \in T$, then we apply the following ‘blindness’ argument. Suppose that $y \in T$. Since the cycle C is properly coloured, $c(yz) \neq c(yy_+)$ or $c(yz) \neq c(yy_-)$. We treat y to be in R if $c(yz) \neq c(yy_+)$, and y to be in S otherwise. We apply a similar treatment when $z \in T$. Hence, we are back in the case when $y, z \in R \cup S$. Therefore, the proof of the claim is completed. $\square$

Claim 3.2. *For each vertex $y \in V(F)$ there exists a vertex $u = u(y) \notin V(F)$ such that either $yu \notin E(G)$ or $c(yu) = c(y)$.*

Proof of claim. If $y \in W$, then $c(y)$ is defined to be $c(xy)$ and so we are done by setting $u(y) = x$. Hence, we may assume $y \in R' \cup S'$.

If $y \in R'$ with $y_- \in N_{(d)}^c(x)$, then xy is not an edge. Set $u(y) = x$ for $y \in R$ with $y_- \in N_{(d)}^c(x)$. If $y \in R'$ with $y_- \in N_{(f)}^c(x)$, then $c(xy_-) = c(xz)$ for $z \in V(C_{y_-})$ by the remark after the definition of $N^c(x)$. Hence, $V(C_y) \cap N^c(x) = \{y_-\}$. This implies that $V(C_y) \cap V(F) = \{y\}$ and so $y_+ \notin V(F)$. Set $u(y) = y_+$ for $y \in R$ with $y_- \in N_{(f)}^c(x)$.

For each remaining vertex $y \in R' \cup S'$, we are going to show that if $xy \in E(G)$ then $c(xy) = c(y)$. Thus, we are done by setting $u(y) = x$. Hence, xy is assumed to be an edge for the remainder of the proof.

Suppose that $y \in R'$ with $y_- \notin N_{(d)}^c(x) \cup N_{(f)}^c(x)$ (i.e. $y_- \in N_{(b)}^c(x) \cup N_{(e)}^c(x)$). If $y_- \in N_{(d)}^c(x)$, then $c(xy) \neq c(xy_-)$ by definition. If $y_- \in N_{(b)}^c(x)$, then $c(xy_-) = c(yy_-)$. Recall that all monochromatic subgraphs of G are a disjoint union of stars. Hence, $c(xy) \neq c(xy_-)$. In summary, we have

$$c(xy) \neq c(xy_-) \quad \text{for } y \in R' \text{ with } y_- \notin N_{(d)}^c(x) \cup N_{(f)}^c(x). \quad (3.4)$$

If $c(xy) \neq c(yy_+)$, then $xyC_y^+y_-x$ is a properly coloured cycle and so we can enlarge H , a contradiction. Hence, $c(xy) = c(yy_+) = c(y)$ for $y \in R'$ with

$y_- \notin N_{(d)}^c(x) \cup N_{(f)}^c(x)$. By a similar argument, we may assume that

$$c(xy) = c(yy_-) \text{ for } y \in S'. \quad (3.5)$$

Recall that we have already dealt with the case when $y \in R'$ and $y_- \notin N_{(d)}^c(x) \cup N_{(f)}^c(x)$ at the beginning of the proof. This means that claim is true for all $y \in R \cup S$.

Suppose that $y \in T = R' \cap S'$. We may further assume that $y_- \notin N_{(d)}^c(x) \cup N_{(f)}^c(x)$. Since $y \in R'$ and $y_- \notin N_{(d)}^c(x) \cup N_{(f)}^c(x)$, $c(xy) = c(yy_+)$ by (3.4). On the other hand, since $y \in S'$, we have $c(xy) = c(yy_-)$ by (3.5). Hence, $c(yy_+) = c(yy_-)$ contradicting the fact that the cycle C_y is properly coloured. Therefore, $xy \notin E(G)$ for all $y \in T$, so we can set $u(y) = x$ for $y \in T$. $\square$

If $V(F) = T$, then $|T| \geq \delta/2$ by (3.1). By Claim 3.1, $G[T]$ is empty. Claim 3.2 implies that for each $y \in T$ there exists a vertex $u(y) \notin V(F) \cup N(y)$ as the colour $c(y) = c_0$ does not appear in G . Therefore

$$\delta \leq d^c(y) \leq |N_G(y)| \leq n - |T| - |u(y)| \leq n - \delta/2 - 1,$$

which gives $\delta < 2n/3$ as required. Thus, we may assume that $V(F) \neq T$. Next, we are going to show that there exists a vertex $y_0 \in R \cup S \cup W$ such that

$$|\{z \in V(F) \setminus \{y_0\} : y_0z \notin E(G) \text{ or } c(y_0z) = c(y_0)\}| \geq (\delta - 1)/2. \quad (3.6)$$

Define F' to be the directed graph on $V(F)$ such that there is a directed edge from y to z if $yz \notin E(F)$ or $c(yz) = c(z)$. Thus, in proving (3.6), it suffices to show that there exists a vertex $y_0 \in R \cup S \cup W$ with in-degree at least $(\delta - 1)/2$ in F' . By Claim 3.1, the base graph of F' is complete. Moreover, $\vec{yz}$ exists for all $y \in T$ and $z \in V(F)$ as the colour $c(y) = c_0$ does not appear in G . Hence, the number of directed edges into $R \cup S \cup W$ is at least $\binom{|R \cup S \cup W|}{2} + |T||R \cup S \cup W|$. By an averaging argument, there exists $y_0 \in R \cup S \cup W$ with in-degree

$$d^-(y_0) \geq (|R| + |S| + |W| - 1)/2 + |T| \stackrel{(3.1)}{\geq} (\delta - 1)/2,$$

so (3.6) holds. Recall that the colour degree $d^c(y_0) \geq \delta^c(G) = \delta$, so y_0 meets at least δ edges of distinct colours. In particular, there are at least $\delta - 1$ neighbours z' of y_0 with $c(y_0z') \neq c(y_0)$. Hence, there are at most $n - \delta$ vertices z in $V \setminus \{y_0\}$ such that either $y_0z \notin E(G)$ or $c(y_0z) = c(y_0)$. Recall Claim 3.2 that there exists $u = u(y_0) \in V \setminus V(F)$ such that $y_0u \notin E(G)$ or $c(y_0u) = c(y_0)$. Together with (3.6), we have

$$\begin{aligned} n - \delta &\geq |\{z \in V(F) \setminus \{y_0\} : y_0z \notin E \text{ or } c(y_0z) = c(y_0)\}| + |u(y_0)| \\ &\geq (\delta - 1)/2 + 1. \end{aligned}$$

Thus, $\delta < 2n/3$ as required. This completes the proof of Theorem 1.2. $\square$

The properly coloured 2-factor obtained by Theorem 1.2 may contain $|G|/3$ cycles. We would like to bound the number of cycles in a properly coloured 2-factor. In the next lemma, we show that this can be achieved by assuming a slightly larger $\delta_1^c(G)$. Recall that $\delta_1^c(G)$ is the minimum $\delta(G - H)$ over all monochromatic subgraphs H in G . So $\delta_1^c(G) \geq \delta^c(G) - 1$.

Lemma 3.3. *For every integer $k \geq 1$, every edge-coloured graph G with $|G| = n$ and $\delta_1^c(G) \geq 2n/3 + k$ contains a properly coloured 2-factor such that every cycle has length at least $k/2$. In particular, G can be covered by at most $2n/k$ vertex-disjoint properly coloured paths.*

The idea of the proof is rather simple. Recall that a 1-path-cycle is a vertex-disjoint union of at most one path P and a number of cycles. We consider a properly coloured 1-path-cycle H in G such that every cycle has length at least $k/2$ and $|H|$ is maximal. Suppose that $V(H) = V(G)$, so we may assume that H contains a path P or else we are done. Our aim is to show that there exists a properly coloured 2-factor H' in $G[V(P)]$ such that each cycle has length at least $k/2$. Therefore $H - P + H'$ is the desired properly coloured 2-factors.

Proof of Lemma 3.3. Suppose the contrary and let G be a counterexample with $|G| = n$ and $\delta_1^c(G) \geq 2n/3 + k$. Let H be a properly coloured 1-path-cycle in G such that every cycle has length at least $k/2$ and $|H|$ is maximal. If $V(H) = V(G)$, then H contains a path P or else we are done. If $V(H) \neq V(G)$, then we may assume H contains a path (maybe consist of a single vertex). Let $P = z_1 z_2 \dots z_\ell$ be the path in H . We further assume that H is chosen such that ℓ maximum (subject to $|H|$ is maximal). Suppose that $\ell = 1$ and so $P = z_1$. If $N(z_1) \setminus V(H) \neq \emptyset$, then we can extend P contradicting the maximality of $|H|$. Hence, we may assume that $y \in V(H) \cap N(z_1)$. Orient C_y into a directed cycle such that $c(yy_+) \neq c(z_1 y)$. Then $H - C_y + z_1 y C_y^+ y_-$ is a properly coloured 1-path-cycle on vertex set $V(H)$ with path $z_1 y C_y^+ y_-$ contradicting the maximality of ℓ . Therefore, $\ell \geq 2$.

Let $N_1 = \{x \in N_G(z_1) : c(z_1 x) \neq c(z_1 z_2)\}$ and $N_\ell = \{x \in N_G(z_\ell) : c(z_\ell x) \neq c(z_\ell z_{\ell-1})\}$. So

$$|N_1|, |N_\ell| \geq \delta_1^c(G) \geq 2n/3 + k. \quad (3.7)$$

By maximality of $|H|$, $N_1, N_\ell \subseteq V(H)$. If $y \in N_1 \setminus V(P)$, then orient C_y into a directed cycle such that $c(yy_+) \neq c(z_1 y)$. Then $H - C_y - P + y_+ C_y^- y P$ contradicts the maximality of ℓ . Therefore, $N_1 \subseteq V(P)$ and similarly $N_\ell \subseteq V(P)$. Moreover,

$$\ell = |V(P)| \geq |N_1| \geq \delta_1^c(G) \geq 2n/3 + k.$$

If $z_1 \in N_\ell$ and $c(z_1 z_\ell) \neq c(z_1 z_2)$, then $C = z_1 \dots z_\ell z_1$ is a properly coloured cycle of length at least $\ell \geq k$. Hence, $H - P + C$ consists of vertex-disjoint properly coloured cycles each of length at least $k/3$, a contradiction. Therefore, if $z_1 \in N_\ell$, then $c(z_1 z_\ell) = c(z_1 z_2)$. A similar statement also holds if $z_\ell \in N_1$. Let

$$N'_q = N_q \setminus (\{z_1, z_2, \dots, z_{\lceil k/2 \rceil}\} \cup \{z_\ell, z_{\ell-1}, \dots, z_{\ell - \lceil k/2 \rceil + 1}\}) \quad (3.8)$$

for $q \in \{1, \ell\}$. Since $z_1 \notin N_1$ and $z_\ell \notin N_\ell$, by (3.7), we have

$$|N'_1|, |N'_\ell| \geq 2n/3 + k - (2\lceil k/2 \rceil - 1) \geq 2n/3. \quad (3.9)$$

For a given $x = z_i \in V(P)$, we write x_+ and x_- for z_{i+1} and z_{i-1} respectively. Set $c_-(x) = c(xx_-)$ and $c_+(x) = c(xx_+)$. For distinct $x, y \in V(P)$, xPy

denotes the subpath of P from x to y . If $x = y$, set $xPx = x$. For $q \in \{1, \ell\}$, define

$$\begin{aligned} R_q &= \{x : x_- \in N'_q \text{ and } c(x_-z_q) = c_+(x_-)\}, \\ S'_q &= \{x : x_+ \in N'_q \text{ and } c(x_+z_q) = c_-(x_+)\}, \\ S''_q &= \{x : x_+ \in N'_q \text{ and } c_+(x_+) \neq c(x_+z_q) \neq c_-(x_+)\}, \\ S_q &= S'_q \cup S''_q. \end{aligned}$$

Note that $S'_q \cap S''_q = \emptyset$ and $z_q \notin R_q \cup S_q$. Since P is properly coloured, for each $x \in N_1$, we have $c(xz_1) \neq c_+(x)$ or $c(xz_1) \neq c_-(x)$. Therefore, the three sets $\{x_- : x \in R_1\}$, $\{x_+ : x \in S_1\}$, $\{x_+ : x \in S''_1\}$ are pairwise disjoint. Moreover, the union of these three sets is precisely N'_1 . Hence, there is a one-to-one correspondence between N'_1 and the multiset $R_1 \cup S_1$, and a similar correspondence between N'_ℓ and the multiset $R_\ell \cup S_\ell$. Hence, $|R_q| + |S_q| = |N'_q|$ for $q \in \{1, \ell\}$. Note that if $z_\ell \in R_1 \cup S_1$, then $z_\ell \in R_1 \setminus S_1$. Similarly if $z_1 \in R_\ell \cup S_\ell$, then $z_1 \in S_\ell \setminus R_\ell$. For $q \in \{1, \ell\}$, we set

$$W_q = ((R_q \cup S_q) \setminus \{z_1, z_\ell\}) \cup \{z_q\}, \quad T_q = R_q \cap S_q, \quad U_q = W_q \setminus T_q.$$

Since $z_2 \notin N'_1$, $z_1 \notin R_1 \cup S_1$ and so $U_1 \neq \emptyset$. Also, $z_\ell \notin R_1 \cap S_1$, so $|W_1| + |T_1| \geq |N'_1|$. Similarly, $U_\ell \neq \emptyset$ and $|W_\ell| + |T_\ell| \geq |N'_\ell|$. Moreover, for $q \in \{1, \ell\}$,

$$2|W_q| = |W_q| + |T_q| + |U_q| \geq |N'_q| + |U_q| \stackrel{(3.9)}{\geq} |U_q| + 2n/3. \quad (3.10)$$

Now define an auxiliary directed bipartite graph F with vertex classes W_1 and W_ℓ such that for $\{q, q'\} = \{1, \ell\}$ there is a directed edge from $x \in W_q$ to $y \in W_{q'}$ if and only if $xy \in E(G)$ and one of the following statements holds:

- (i) $x \in R_q \setminus (T_q \cup \{z_1, z_\ell\})$ and $c(xy) \neq c_+(x)$;
- (ii) $x \in S_q \setminus (T_q \cup \{z_1, z_\ell\})$ and $c(xy) \neq c_-(x)$;
- (iii) $x \in T_q$;
- (iv) $q = 1$, $x = z_1$ and $c(xy) \neq c(z_1z_2)$;
- (v) $q = \ell$, $x = z_\ell$ and $c(xy) \neq c(z_\ell z_{\ell-1})$.

We are going to show that there exists a vertex $u \in U_1 \cup U_\ell$ such that u is in at least k directed 2-cycles in F . If $x \in R_1 \setminus T_1$, then $\overrightarrow{xy}$ exists of $y \in W_\ell$ if and only if $c(xy) \neq c_+(x)$. Hence, the out-degree of x in F is

$$\begin{aligned} d_F^+(x) &\geq |\{z \in N(x) : c(xz) \neq c_+(x)\}| + |W_\ell| - n \\ &\geq \delta_1^c(G) + |W_\ell| - n \\ &\stackrel{(3.10)}{\geq} (2n/3 + k) + \frac{|U_q| + 2n/3}{2} - n = (|U_\ell| + 2k)/2. \end{aligned}$$

A similar statement holds for $x \in (S_1 \setminus T_1) \cup z_1$. Recall that $U_1 = (R_1 \cup S_1 \cup z_1) \setminus T_1$. In summary, for all $x \in U_1$,

$$d_F^+(x) \geq (|U_\ell| + 2k)/2.$$

Hence, there are at least $|U_1|(|U_\ell| + 2k)/2$ directed edges from U_1 to W_ℓ , and similarly there are at least $|U_\ell|(|U_1| + 2k)/2$ directed edges from U_ℓ to W_1 . Note that if $\overrightarrow{xy}$ is a directed edge in F with $x \in U_1$ and $y \in T_\ell$, then $xy \in E(G)$ and so $\{x, y\}$ forms a directed 2-cycle in F by (iii). Hence, if $\overrightarrow{xy}$ is a directed edge in F not contained in a directed 2-cycle with $x \in U_1$

and $y \in W_\ell$, then $y \in U_\ell$. A similar statement holds for $\overrightarrow{xy}$ with $x \in U_\ell$ and $y \in W_1$. Therefore, at most $|U_1||U_\ell|$ directed edges from $U_1 \cup U_\ell$ are not contained in a directed 2-cycle in F . The number of directed edges $\overrightarrow{xy}$ in F such that $x \in U_1 \cup U_\ell$ and $\overrightarrow{xy}$ is contained in a directed 2-cycle is at least

$$\frac{|U_1|(|U_\ell| + 2k)}{2} + \frac{|U_\ell|(|U_1| + 2k)}{2} - |U_1||U_\ell| = k(|U_1| + |U_\ell|).$$

Hence, by an averaging argument, there exists a vertex $u \in U_1 \cup U_\ell$ that is in at least k directed 2-cycles in F . Assume that $u \in U_1$. There are at least k vertices $w \in W_2 \setminus u$ such that uw is a directed 2-cycle in F . Pick one such $w \in W_2$ such that the subpath uPw in G has length at least $k/2$. Similarly, if $u \in U_\ell$, then there is a vertex $w \in W_1$ such that uw is a directed 2-cycle in F and the subpath uPw in G has length at least $k/2$. Without loss of generality, we may assume that $u \in W_1$ and $w \in W_\ell$.

If $u \in T_1 = R_1 \cap S_1$, then $u \in R_1 \setminus \{z_1, z_\ell\}$ with $c(uw) \neq c_+(u)$ or $u \in S_1 \setminus \{z_1, z_\ell\}$ with $c(uw) \neq c_-(u)$ as P is a properly coloured path and $u \notin \{z_1, z_\ell\}$. Together with (i), (ii) and (iv), we may assume that at least one of the following statements hold:

- (a'_1) $u \in R_1$ and $c(uw) \neq c_+(u)$;
- (a'_2) $u \in S_1$ and $c(uw) \neq c_-(u)$;
- (a'_3) $u = z_1$ and $c(z_1w) = c(uw) \neq c(z_1z_2)$.

If $u \in R_1$, then the definition of R_1 implies that $u_- \in N'_1$ and $c(z_1u_-) = c_+(u_-) \neq c_-(u_-)$. Since $u_- \in N'_1$, $c(z_1z_2) \neq c(z_1u_-)$. A similar statement also holds for $u \in S_1$, so at least one of the following statements hold:

- (a_1) $u \in R_1$, $c(uw) \neq c_+(u)$ and $c(z_1z_2) \neq c(z_1u_-) \neq c_-(u_-)$;
- (a_2) $u \in S_1$, $c(uw) \neq c_-(u)$ and $c(z_1z_2) \neq c(z_1u_+) \neq c_+(u_+)$;
- (a_3) $u = z_1$ and $c(z_1w) \neq c(z_1z_2)$.

By symmetry, we can deduce that at least one of the following statements hold for w


- (b_1) $w \in R_\ell$, $c(uw) \neq c_+(w)$ and $c(z_\ell z_{\ell-1}) \neq c(z_\ell w_-) \neq c_-(w_-)$;
- (b_2) $w \in S_\ell$, $c(uw) \neq c_-(w)$ and $c(z_\ell z_{\ell-1}) \neq c(z_\ell w_+) \neq c_+(w_+)$;
- (b_3) $w = z_\ell$ and $c(z_\ell u) \neq c(z_{\ell-1}z_\ell)$.

We now claim that

- (†) there exists a properly coloured 2-factor H' in $G[V(P)]$ such that each cycle has length at least $k/2$.

If (†) holds, then $H'' = H - P + H'$ is a disjoint union of properly coloured cycles each of length at least $k/2$ on vertex set $V(H)$. If $V(H'') = V(G)$, then H'' is the desired properly coloured 2-factor. If $z \in V(G) \setminus V(H'')$, then $H'' \cup z$ is a properly coloured 1-path-cycle contradicting the maximality of $|H| = |H''|$. Therefore, in order to complete the proof, it suffices to prove (†) for each combination of (a_i) and (b_j) for $1 \leq i, j \leq 3$. We say that $u < w$ if $u = z_i$ and $w = z_j$ with $i < j$. We would like to point out that Cases 2–6 are proved by similar arguments used in Case 1. Since the explicit structures of H' (see Figure 2) are different, we include their proofs for completeness.

Case 0: (a_3) and (b_3) hold. Recall that if $z_1 \in N_\ell$, then $c(z_1z_\ell) = c(z_1z_2)$. Hence, at least one of (a_3) and (b_3) does not hold, a contradiction.

FIGURE 2. Structures of H'

Case 1: (a₁) and (b₂) hold. We have the following three statements:

$$c_+(u) \neq c(uw) \neq c_-(w), \quad (3.11)$$

$$c(z_1 z_2) \neq c(z_1 u_-) \neq c_-(u_-), \quad (3.12)$$

$$c(z_\ell z_{\ell-1}) \neq c(z_\ell w_+) \neq c_+(w_+). \quad (3.13)$$

If $u_- = w$, then we set $H' = z_1 P u_- z_1 + w_+ P z_\ell w_+$ (see Figure 2 (A)). Note that H' is properly coloured and a union of two cycles on vertex set $V(H)$. By (a₁) and (3.8), we have $u_- \in N'_1 \subseteq N_1 \setminus \{z_1, \dots, z_{\lceil k/2 \rceil}\}$. Thus, $z_1 P u_- z_1$ is a cycle of length at least $k/2$. By a similar argument, we deduce that $w_+ P z_\ell w_+$ is also a cycle of length at least $k/2$ as $w_+ \in N'_\ell$ by (b₂). Hence (†) holds.

Next suppose that $u_- > w$. If $u_- = w_+$ and $c(z_\ell w_+) = c(z_1 u_-)$, then since $u \in R_1$,

$$c(z_\ell w_+) = c(z_1 u_-) = c_+(u_-) = c_+(w_+) \neq c_-(w_+).$$

This contradicts the fact that $w \in S_\ell$. Therefore, if $u_- = w_+$, then $c(z_1 u_-) \neq c(z_\ell w_+)$. Together with (3.11)–(3.13), we conclude that $H' = z_1 u_- P w_+ z_1 P u w P z_1$ (see Figure 2 (B)) is a properly coloured spanning cycle on $V(P)$, so (†) holds.

If $u < w$, then $z_1 P u_- z_1$, $u P w u$ and $w_+ P z_\ell w_+$ (see Figure 2 (C)) are properly coloured cycles by (3.11)–(3.13). We will show that each is a cycle of length at least $k/2$ (which then implies (†)). By (a₁) and (3.8), we have $u_- \in N'_1 \subseteq N_1 \setminus \{z_1, \dots, z_{\lceil k/2 \rceil}\}$. Thus, $z_1 P u_- z_1$ is a cycle of length at least $k/2$. Similarly, $w_+ P z_\ell w_+$ is also a cycle of length at least $k/3$ as $w_+ \in N'_\ell$ by (b₂). Since u and w are chosen such that the subpath $u P w$ has length at least $k/2$, $u P w u$ has length at least $k/2$. Therefore, (†) holds for (a₁) and (b₂).

Case 2: (a₁) and (b₁) hold. We have the following three statements:

$$c_+(u) \neq c(uw) \neq c_+(w), \quad (3.14)$$

$$c(z_1 z_2) \neq c(z_1 u_-) \neq c_-(u_-), \quad (3.15)$$

$$c(z_\ell z_{\ell-1}) \neq c(z_\ell w_-) \neq c_-(w_-). \quad (3.16)$$

If $u < w$, then we set $H' = z_1 P u_- z_1 + u P w_- z_\ell P w u$ (see Figure 2 (D)). Note that H' is properly coloured. By (a₁) and (3.8), we have $u_- \in N'_1 \subseteq N_1 \setminus \{z_1, \dots, z_{\lceil k/2 \rceil}\}$. Thus, $z_1 P u_- z_1$ is a cycle of length at least $k/2$. Recall our choices of u and w that $u P w$ has length at least $k/2$. This implies that $u P w_- z_\ell P w u$ is also a cycle of length at least $k/2$. Hence (†) holds.

Next suppose that $u > w$. By (3.14), we deduce that $w_+ \neq u$ and so $w < u_-$. Hence, $w P u_-$ is a path with length at least one. Together with (3.14)–(3.16), we conclude that $H' = z_1 P w_- z_\ell P u w P u_- z_1$ (see Figure 2 (E)) is a properly coloured spanning cycle on $V(P)$, so (†) holds. Therefore, (†) holds for (a₁) and (b₁).

Case 3: (a₁) and (b₃) hold. Since $w = z_\ell$, we have

$$c_+(u) \neq c(u z_\ell) \neq c_-(z_{\ell-1} z_\ell) \text{ and } c(z_1 z_2) \neq c(z_1 u_-) \neq c_-(u_-). \quad (3.17)$$

Note that both $z_1 P u_- z_1$ and $u P z_\ell u$ (see Figure 2 (F)) are properly coloured cycles. By (3.8), both cycles have length at least $k/2$. Therefore, (†) holds for (a₁) and (b₃) by setting $H' = z_1 P u_- z_1 + u P z_\ell u$.

Case 4: (a₂) and (b₁) hold. We have the following three statements:

$$c_-(u) \neq c(uw) \neq c_+(w), \quad (3.18)$$

$$c(z_1z_2) \neq c(z_1u_+) \neq c_+(u_+), \quad (3.19)$$

$$c(z_\ell z_{\ell-1}) \neq c(z_\ell w_-) \neq c_-(w_-). \quad (3.20)$$

First suppose that $u < w_-$. If $u_+ = w_-$ and $c(z_1u_+) = c(z_\ell w_-)$, then since $w \in R_\ell$,

$$c(z_1u_+) = c(z_\ell w_-) = c_+(w_-) = c_+(u_+) \neq c_-(u_+)$$

This contradicts the fact that $u \in S_1$. Therefore, if $u_+ = w_-$, then $c(z_1u_-) \neq c(z_\ell w_+)$. Together with (3.18)–(3.20), we conclude that $H' = z_1Pu_wPz_\ell w_-Pu_+z_1$ (see Figure 2 (G)) is a properly coloured spanning cycle on $V(P)$, so $(\dagger)$ holds.

If $u = w_-$, then $H' = z_1Pw_-z_\ell Pu_+z_1$ (see Figure 2 (H)) is a properly coloured spanning cycle on $V(P)$ by (3.19) and (3.20). Thus $(\dagger)$ holds.

Suppose that $u > w$. By (3.18), we deduce that $w_+ \neq u$ and so $w < u_-$. Hence, wPu_w is a properly coloured cycle. Moreover, it has length at least $k/2$ by our choices of u and w . By (3.19) and (3.20), we conclude that $z_1Pw_-z_\ell u_+z_1$ is a properly coloured cycle. It has length at least k by (3.8). Hence $(\dagger)$ holds by setting $H' = wPu_w + z_1Pw_-z_\ell Pu_wPu_-z_1$ (see Figure 2 (I)). Therefore, $(\dagger)$ holds for (a₂) and (b₁).

Case 5: (a₂) and (b₃) hold. Since $w = z_\ell$, we have

$$c_-(u) \neq c(uz_\ell) \neq c_-(z_{\ell-1}z_\ell) \text{ and } c(z_1z_2) \neq c(z_1u_+) \neq c_+(u_+).$$

Note that $z_1Pu_zz_\ell Pu_+z_1$ (see Figure 2 (J)) is a properly coloured cycle spanning $V(P)$ and so $(\dagger)$ holds.

Case 6: (a_i) and (b_j) hold for $(i, j) \in \{(3, 2), (2, 2), (3, 1)\}$. Let $P' = z'_1z'_2 \dots z'_\ell$ be the properly coloured path obtained by reversing P . Hence, $z'_i = z_{\ell-i+1}$ for all $1 \leq i \leq \ell$. Given $x' = z'_i \in V(P')$ we abuse the notation and write x_{+}' and x_{-}' to mean z'_{i+1} and z'_{i-1} . Also, let $c_{+}'(x)$ and $c_{-}'(x)$ to mean the colours $c(xx_{+}')$ and $c(xx_{-}')$ respectively. (Hence, these primed notations are defined respect to P' .) Recall that P' is the reversed P , so $c_{+}'(x) = c_-(x)$ and $c_{-}'(x) = c_+(x)$. Further, set $u' = w$ and $w' = u$.

Now suppose that (a₃) and (b₂) hold. Under the primed notation, we have

$$c_{+}'(u') \neq c(u'z'_\ell) \neq c_-(z'_{\ell-1}z'_\ell) \text{ and } c(z'_1z'_2) \neq c(z'_1u_{-}') \neq c_{-}'(u_{-}').$$

More importantly, if we ignore the primes, then we obtain (3.17). Therefore, in hindsight, we are in case when (a₁) and (b₃) hold with respect to P' , i.e. we are in Case 3. Hence, $(\dagger)$ holds by setting $H' = z'_1P'u_{-}'z'_1 + u'P'z'_\ell u'$.

Similarly, we deduced that the case when (a₂) and (b₂) hold with respect to P , corresponds to the case when the case when (a₁) and (b₁) holds with respect to P' . Also, (a₃) and (b₁) (with respect to P) corresponds to (a₂) and (b₃) (with respect to P'). Therefore, we are in Case 1 and Case 5 respectively. The proof of the lemma is completed. $\square$

4. ABSORBING CYCLE

The aim of this section is to prove the following lemma, that is, to show that there exists a small absorbing cycle.

Lemma 4.1 (Absorbing cycle lemma). *Let $0 < \varepsilon < 2^{-9}3^{-1}$. Then exists an integer n_0 such that whenever $n \geq n_0$ the following holds. Suppose that G is an edge-coloured graph of order n with $\delta_1^c(G) \geq (2/3 + \varepsilon)n$. Then there exists a properly coloured cycle C of length at most $2\varepsilon n/3$ such that for all $k \leq (8\varepsilon^2 n)/243$ and for any vertex-disjoint properly coloured paths $P_1, \dots, P_k$ in $G \setminus C$, there exists a properly coloured cycle spanning the vertex set $V(C) \cup \bigcup_{1 \leq i \leq k} V(P_i)$.*

Given a vertex x , we say that a path P is an *absorbing path* for x if the following conditions hold:

- (i) $P = z_1 z_2 z_3 z_4$ is a properly coloured path of length 3;
- (ii) $V(P) \cap \{x\} = \emptyset$;
- (iii) $z_1 z_2 x z_3 z_4$ is a properly coloured path;

Next we define an absorbing path for two disjoint edges. Given two vertex-disjoint edges $x_1 x_2, y_1 y_2$, we say that a path P is an *absorbing path* for $(x_1, x_2; y_2, y_1)$ if the following conditions hold:

- (i) $P = z_1 z_2 z_3 z_4$ is a properly coloured path of length 3;
- (ii) $V(P) \cap \{x_1, x_2, y_1, y_2\} = \emptyset$;
- (iii) both $z_1 z_2 x_1 x_2$ and $y_2 y_1 z_3 z_4$ are properly coloured paths of length 3.

Note that the ordering of $(x_1, x_2; y_2, y_1)$ is important. Given a vertex x , let $\mathcal{L}(x)$ be the set of absorbing paths for x . Similarly, given two vertex-disjoint edges $x_1 x_2, y_1 y_2$, let $\mathcal{L}(x_1, x_2; y_2, y_1)$ be the set of absorbing paths for $(x_1, x_2; y_2, y_1)$. By the definition of an absorbing path for $(x_1, x_2; y_2, y_1)$, we obtain the following simple proposition.

Proposition 4.2. *Let $P' = x_1 x_2 \dots x_{\ell-1} x_\ell$ be a properly coloured path with $\ell \geq 4$. Let $P = z_1 z_2 z_3 z_4$ be an absorbing path for $(x_1, x_2; x_{\ell-1}, x_\ell)$ with $V(P) \cap V(P') = \emptyset$. Then $z_1 z_2 x_1 x_2 \dots x_{\ell-1} x_\ell z_3 z_4$ is a properly coloured path.*

Lemma 4.1 will be proved as follows. Suppose that $\delta_1^c(G) \geq (2/3 + \varepsilon)n$. In the next lemma, Lemma 4.3, we show that every $\mathcal{L}(x)$ and every $\mathcal{L}(x_1, x_2; y_2, y_1)$ are large. By a simple probabilistic argument, Lemma 4.4 shows that there exists a small family $\mathcal{F}'$ of vertex-disjoint properly coloured paths (of order 4) such that $\mathcal{F}'$ contains at a linear number of members of $\mathcal{L}(x)$ for all $x \in V(G)$ and a similar statement for $\mathcal{L}(x_1, x_2; y_2, y_1)$. Finally, we join all paths in $\mathcal{F}'$ into one short properly coloured cycle C using Lemma 4.5. Moreover, C satisfies the desired property in Lemma 4.1.

Lemma 4.3. *Let $0 < \varepsilon < 1/2$ and let $n \geq \varepsilon^{-2}$ be an integer. Suppose that G is an edge-coloured graph on n vertices with $\delta_1^c(G) \geq (1/2 + \varepsilon)n$. Then $|\mathcal{L}(x)| \geq \varepsilon^3 n^4$ for every $x \in V(G)$ and $|\mathcal{L}(x_1, x_2; y_2, y_1)| \geq \varepsilon^3 n^4$ for every four distinct vertices $x_1, x_2, y_1, y_2 \in V(G)$ with $x_1 x_2, y_1 y_2 \in E(G)$.*

Proof. Fix a vertex $x \in V(G)$. Choose a vertex $z_2 \in N(x)$. Next pick another vertex $z_3 \in N(z_2) \cap N(x)$ such that $c(xz_2) \neq c(z_2 z_3) \neq c(xz_3)$. Notice that the number of such z_3 is at least $2\delta_1^c(G) - n \geq 2\varepsilon n$. Since $\delta_1^c(G) \geq (1/2 + \varepsilon)n$, $\Delta(H) \leq (1/2 - \varepsilon)n$ for all monochromatic subgraphs H in G . Hence, the number of $z_1 \in V(G) \setminus \{x, z_2, z_3\}$ such that $c(z_2 z_3) \neq c(z_1 z_2) \neq c(z_2 x)$ is at least $\delta_1^c(G) - (1/2 - \varepsilon)n - 2 \geq \varepsilon n$. Fix one such z_1 . Similarly, there are at least εn choices for $z_4 \in V(G) \setminus \{x, z_1, z_2, z_3\}$ such that $c(z_2 z_3) \neq c(z_3 z_4) \neq c(xz_3)$. Notice that $z_1 z_2 z_3 z_4$ is an absorbing path

for x . Recall that every path is assumed to be directed. Therefore, there are at least $\delta_1^c(G) \times 2\epsilon n \times \epsilon n \times \epsilon n \geq \epsilon^3 n^4$ many absorbing paths for x .

Next, fix vertex-disjoint edges x_1x_2 and y_1y_2 in G . Choose a vertex $z_2 \in N(x_1) \setminus \{x_1, x_2, y_1, y_2\}$ such that $c(x_1z_2) \neq c(x_1x_2)$. Pick another vertex $z_3 \in N(z_2) \setminus \{x_1, x_2, y_1, y_2\}$ such that $c(xz_2) \neq c(z_2z_3)$ and $c(y_1z_3) \neq c(y_1y_2)$. The number of such z_3 is at least $2\delta_1^c(G) - n - 1 \geq 2\epsilon n - 1$. We then extend z_2z_3 into an absorbing path for $(x_1, x_2; y_2, y_1)$ by a similar construction used above. Furthermore, there are at least $(\delta_1^c(G) - 2) \times (2\epsilon n - 1) \times \epsilon n \times \epsilon n \geq \epsilon^3 n^4$ many absorbing paths for $(x_1, x_2; y_2, y_1)$. $\square$

The next lemma is proved by a simple probabilistic argument since each of $\mathcal{L}(x_1, x_2; y_2, y_1)$ and $\mathcal{L}(x_1, x_2; y_2, y_1)$ is large.

Lemma 4.4. *Let $0 < \gamma < 1$. Then exists an integer n_0 such that whenever $n \geq n_0$ the following holds. Let G be an edge-coloured graph on n vertices. Suppose that $|\mathcal{L}(x)| \geq \gamma n^4$ for every $x \in V(G)$ and $|\mathcal{L}(x_1, x_2; y_2, y_1)| \geq \gamma n^4$ for every four distinct vertices $x_1, x_2, y_1, y_2 \in V(G)$ with $x_1x_2, y_1y_2 \in E(G)$. Then there exists a family $\mathcal{F}'$ of vertex-disjoint properly coloured paths each of length 3 such that*

$$\begin{aligned} |\mathcal{F}'| &\leq 2^{-6}\gamma n, \\ |\mathcal{L}(x) \cap \mathcal{F}'| &\geq 2^{-9}\gamma^2 n \quad \text{for all } x \in V(G), \\ |\mathcal{L}(x_1, x_2; y_2, y_1) \cap \mathcal{F}'| &\geq 2^{-9}\gamma^2 n \end{aligned}$$

for every four distinct vertices $x_1, x_2, y_1, y_2 \in V(G)$ with $x_1x_2, y_1y_2 \in E(G)$.

Proof. Recall that each path is assumed to be directed. So a path $z_1z_2z_3z_4$ will be considered as a 4-tuple (z_1, z_2, z_3, z_4) . Choose a family $\mathcal{F}$ of 4-tuples in $V(G)$ by selecting each of the $n!/(n-4)!$ possible 4-tuples independently at random with probability

$$p = 2^{-7}\gamma \frac{(n-4)!}{(n-1)!} \geq 2^{-7}\gamma n^{-3}.$$

Then by Chernoff's bound (see e.g. [2]) with probability $1 - o(1)$ as $n \rightarrow \infty$, the family $\mathcal{F}$ satisfies the following properties:

$$|\mathcal{F}| \leq 2p \frac{n!}{(n-4)!} = 2^{-6}\gamma n \quad (4.1)$$

and, for every $x \in V(G)$,

$$|\mathcal{L}(x) \cap \mathcal{F}| > \frac{p}{2} |\mathcal{L}(x_1, x_2; y_2, y_1)| \geq 2^{-8}\gamma^2 n \quad (4.2)$$

and, for every four distinct $x_1, x_2, y_1, y_2 \in V(G)$ with $x_1x_2, y_1y_2 \in E(G)$,

$$|\mathcal{L}(x_1, x_2; y_2, y_1) \cap \mathcal{F}| > \frac{p}{2} |\mathcal{L}(x_1, x_2; y_2, y_1)| \geq 2^{-8}\gamma^2 n. \quad (4.3)$$

We say that two 4-tuples (a_1, a_2, a_3, a_4) and (b_1, b_2, b_3, b_4) are *intersecting* if $a_i = b_j$ for some $1 \leq i, j \leq 4$. Furthermore, we can bound the expected number of intersecting 4-tuples in $\mathcal{F}$ from above by

$$\frac{n!}{(n-4)!} \times 4^2 \times \frac{(n-1)!}{(n-4)!} \times p^2 = 2^{-10}\gamma^2 n.$$

Thus, using Markov's inequality, we derive that with probability at least $1/2$,

$$\mathcal{F} \text{ contains at most } 2^{-9}\gamma^2n \text{ intersecting pairs.} \quad (4.4)$$

Hence, with positive probability the family $\mathcal{F}$ has all properties stated in (4.1)–(4.4). By deleting all the intersecting 4-tuples and those 4-tuples not absorbing paths in such a family $\mathcal{F}$, we get a subfamily $\mathcal{F}'$ consisting of pairwise disjoint 4-tuples, which satisfies

$$|\mathcal{L}(x) \cap \mathcal{F}'| > 2^{-8}\gamma^2n - 2^{-9}\gamma^2n = 2^{-9}\gamma^2n$$

for every $x \in V(G)$ and a similar statement holds for $|\mathcal{L}(x_1, x_2; y_2, y_1) \cap \mathcal{F}'|$. Since each 4-tuple in $\mathcal{F}'$ is an absorbing path, $\mathcal{F}'$ is a set of vertex-disjoint properly coloured paths of length 3. $\square$

In order to prove Lemma 4.1, it is sufficient to join the paths in $\mathcal{F}'$ given by Lemma 4.4 into a short properly coloured cycle C . The next lemma shows that we can join any two disjoint edges into a properly coloured path of length 5.

Lemma 4.5. *Let $n, k \geq 1$ be integers. Suppose that G is an edge-coloured graph of order n with $\delta_1^c(G) \geq 2n/3 + k$. Let x_1x_2, y_1y_2 be two edges in G with $x_2 \neq y_2$. Then there exist $(3k-2)(2n/3 + k - 2)$ paths z_1z_2 with $z_1, z_2 \in V(G) \setminus \{x_1, x_2, y_1, y_2\}$ such that both $x_1x_2z_1z_2$ and $z_1z_2y_2y_1$ are properly coloured paths. In particular, if x_1, x_2, y_1 and y_2 are distinct, then $x_1x_2z_1z_2y_2y_1$ is a properly coloured path.*

Proof. Let X be the set of vertices $x \in N(x_2) \setminus \{x_1, x_2, y_1, y_2\}$ such that $c(xx_2) \neq c(x_1x_2)$. Similarly, let Y be the set of vertices $y \in N(y_2) \setminus \{x_1, x_2, y_1, y_2\}$ such that $c(yy_2) \neq c(y_1y_2)$. So $|X|, |Y| \geq \delta_1^c(G) - 2 \geq 2n/3 + k - 2$. Define an auxiliary bipartite directed graph H with vertex classes X and Y such that for $x \in X$ and $y \in Y$

- (a) $\vec{xy} \in E(H)$ if and only if $xy \in E(G)$ and $c(xy) \neq c(xx_2)$;
- (b) $\vec{yx} \in E(H)$ if and only if $xy \in E(G)$ and $c(xy) \neq c(yy_2)$.

The outdegree of each $x \in X$ in H is

$$\begin{aligned} d^+(x) &= |Y \cap \{z \in N_G(x) : c(xz) \neq c(xx_2)\}| \\ &\geq |Y| + \delta_1^c(G) - n \geq (|Y| + 3k - 2)/2 \end{aligned}$$

and similarly for each $y \in Y$, $d^+(y) \geq (|X| + 3k - 2)/2$. Hence the number of directed 2-cycles in H is at least

$$\begin{aligned} &\sum_{x \in X} d^+(x) + \sum_{y \in Y} d^+(y) - |X||Y| \\ &\geq \frac{|X|(|Y| + 3k - 2)}{2} + \frac{|Y|(|X| + 3k - 2)}{2} - |X||Y| \\ &= \frac{(|X| + |Y|)(3k - 2)}{2} \geq (3k - 2)(2n/3 + k - 2). \end{aligned}$$

Let z_1z_2 be a directed 2-cycle in H with $z_1 \in X$ and $z_2 \in Y$. This implies that $c(x_2z_1) \neq c(z_1z_2) \neq c(z_2y_2)$. Hence, $x_1x_2z_1z_2$ and $z_1z_2y_2y_1$ are properly coloured paths. Therefore, the lemma follows. $\square$

We are ready to prove Lemma 4.1.

Proof of Lemma 4.1. Let $\gamma = 2^6\varepsilon/9$. By Lemma 4.3 (taking $\varepsilon = 1/6$), $\mathcal{L}(x) \geq \gamma n^4$ for all $x \in V(G)$ and $\mathcal{L}(x_1, x_2; y_2, y_1) \geq \gamma n^4$ for every four distinct vertices $x_1, x_2, y_1, y_2 \in V(G)$ with $x_1x_2, y_1y_2 \in E(G)$. Let $\mathcal{F}'$ be the set of properly coloured paths obtained by Lemma 4.4. Therefore, $|\mathcal{F}'| \leq 2^{-6}\gamma n = \varepsilon n/9$ and

$$|\mathcal{L}(x) \cap \mathcal{F}'| \geq 2^{-9}\gamma^2 n = \frac{8\varepsilon^2 n}{81}, \quad |\mathcal{L}(x_1, x_2; y_2, y_1) \cap \mathcal{F}'| \geq \frac{8\varepsilon^2 n}{81} \quad (4.5)$$

for all $x \in V(G)$ and every four distinct vertices $x_1, x_2, y_1, y_2 \in V(G)$ with $x_1x_2, y_1y_2 \in E(G)$. Suppose that $\mathcal{P}$ is a family of at most $(8\varepsilon^2 n)/243$ vertex-disjoint properly coloured paths in $V \setminus V(\mathcal{F}')$. Let $\mathcal{P}'$ be the family of vertex-disjoint properly coloured paths in $V \setminus V(\mathcal{F}')$ obtained from $\mathcal{P}$ by breaking every path $P \in \mathcal{P}$ with $|P| \leq 3$ up into isolated vertices. Hence, $\mathcal{P}'$ contains at most $(8\varepsilon^2 n)/81$ paths and for each path P in $\mathcal{P}'$ either $|P| = 1$ or $|P| \geq 4$. Now, we assign each path P in $\mathcal{P}'$ to a distinct path Q in $\mathcal{F}'$ such that $Q \in \mathcal{L}(V(P))$ if $|P| = 1$ and $Q \in \mathcal{L}(x_1, x_2; x_{\ell-1}, x_\ell)$ if $P = x_1x_2 \dots x_\ell$ and $\ell \geq 4$. The assignment exists by (4.5). Apply Proposition 4.2 to each pair (P, Q) for P in $\mathcal{P}'$ and obtain a family $\mathcal{F}''$ for vertex-disjoint paths such that $|\mathcal{F}'| = |\mathcal{F}''|$. Note that

$$V(\mathcal{F}'') = V(\mathcal{F}') \cup V(\mathcal{P}') = V(\mathcal{F}') \cup V(\mathcal{P}).$$

Moreover, there is an one-to-one correspondence between each path $P' \in \mathcal{F}'$ and $P'' \in \mathcal{F}''$ such that the endedges of P' and P'' are the same. Therefore, in order to prove Lemma 4.1, it is sufficient to join the paths of $\mathcal{F}'$ into a properly coloured cycle C with $|C| \leq 2\varepsilon n/3$.

Let $P_1, \dots, P_{|\mathcal{F}'|}$ be the properly coloured paths in $\mathcal{F}'$. For each $1 \leq j \leq |\mathcal{F}'|$, we are going to find an edge $e_j = z_1^j z_2^j$ such that $V(e_j) \cap V(\mathcal{F}') = \emptyset$, $P_j z_1^j z_2^j P_{j+1}$ is a properly coloured path and $V(e_j) \cap V(e_{j'}) = \emptyset$ for $j \neq j'$, where we take $P_{|\mathcal{F}'|+1} = P_1$. Assume that we have already found $e_1, \dots, e_{j-1}$. Let $P_j = v_1 v_2 v_3 v_4$ and $P_{j+1} = v'_1 v'_2 v'_3 v'_4$. By Lemma 4.5 taking $x_1 = v_3, x_2 = v_4, y_1 = v'_2$ and $y_2 = v'_1$, there exist at least $4\varepsilon n^2/3$ edges $e = z_1 z_2$ such that $v_3 v_4 z_1 z_2 v'_1 v'_2$ is a properly coloured path. Set $W_j = V(\mathcal{F}') \cup \bigcup_{1 \leq j' < j} V(e_{j'})$, so

$$|W_j| \leq |V(\mathcal{F}')| + 2(j-1) < 4|\mathcal{F}'| + 2|\mathcal{F}'| = 6|\mathcal{F}'| \leq 2\varepsilon n/3.$$

Moreover, the number of edges $z_1 z_2$ incident with $V(Q) \cap V(W_j)$ is at most $|W_j|n < 2\varepsilon n^2/3$. Therefore, there exists an edge $z_1^j z_2^j$ in $G[V \setminus W_j]$ such that $v_3 v_4 z_1^j z_2^j v'_1 v'_2$ is a properly coloured path, which implies that $P_j z_1^j z_2^j P_{j+1}$ is a properly coloured path. This means that there exists a properly coloured cycle C obtained by concatenating $P_1, e_1, P_2, e_2, \dots, P_{|\mathcal{F}'|}, e_{|\mathcal{F}'|}$. Note that $|C| \leq 6|\mathcal{F}'| \leq 2\varepsilon n/3$. This completes the proof of Lemma 4.1. $\square$

5. PROOF OF THEOREM 1.4

First, we prove that G contains a properly coloured triangle if $\delta_1^c(G) > 2n/3 - 1$.

Proposition 5.1. *Let G be an edge-coloured graph on n vertices with $\delta_1^c(G) > 2n/3 - 1$. Then every vertex in G is contained in a properly coloured triangle.*

Proof. Let x be a vertex in G and let c be the edge-colouring on G . Define H to be the directed graph on $N(x)$ with directed edges $\overrightarrow{yz}$ if and only if yz is an edge in $G[N(x)]$ with $c(yz) \neq c(xy) \neq c(xz)$. For $y \in V(H)$, the outdegree of y in H is

$$\begin{aligned} d^+(y) &\geq |\{z \in N(y) : c(yz) \neq c(xy)\}| + |\{z \in N(x) : c(xz) \neq c(xy)\}| - (n-2) \\ &\geq 2\delta_1^c(G) - n + 2. \end{aligned}$$

Note that if yz is a directed 2-cycle in H , then $\{x, y, z\}$ forms a properly coloured triangle in G . Hence, we may assume that H does not contain any directed 2-cycles. Moreover, we may assume that if $\overrightarrow{zy}$ is in H , then $c(zy) = c(xy)$. By an averaging argument, there exists a vertex $y \in V(H)$ such that

$$d^-(y) \geq d^+(y) \geq 2\delta_1^c(G) - n + 2.$$

Therefore

$$\begin{aligned} d_G(y) &\geq |\{z \in N_G(y) : c(yz) = c(xy)\}| + |\{z \in N_G(y) : c(yz) \neq c(xy)\}| \\ &\geq d^-(y) + \delta_1^c(G) \geq 3\delta_1^c(G) - n + 2. \end{aligned}$$

Since $d_G(y) \leq n-1$, the inequality above implies that $\delta_1^c(G) \leq 2n/3 - 1$, a contradiction. $\square$

Finally, we are going to prove Theorem 1.4.

Proof of Theorem 1.4. Let G be an edge-coloured graph on n vertices with $\delta_1^c(G) \geq (2/3 + \varepsilon)n$ as stated in Theorem 1.4. By Proposition 5.1, G contains a properly coloured triangle.

Suppose that ℓ is an integer with $4 \leq \ell \leq \varepsilon n/3$. Since $\delta_1^c(G) \geq (2/3 + \varepsilon)n$, we can greedily constructed a properly coloured path of length $2n/3$. In particular, G contains a properly coloured path $P = x_1x_2 \dots x_{\ell-2}$ of length $\ell - 3$. Let G' be the subgraph of G obtained after removing all the vertices $x_3, x_4, \dots, x_{\ell-4}$. Note that $\delta_1^c(G') \geq 2(n + \varepsilon)/3 \geq 2|G'|/3 + 1$. Hence, Lemma 4.5 implies that there exist distinct $z_1, z_2 \in V(G')$ such that both $x_{\ell-3}x_{\ell-2}z_1z_2$ and $z_1z_2x_1x_2$ are properly coloured paths. Therefore, $x_1x_2 \dots, x_{\ell-2}z_1z_2x_1$ is a properly coloured cycle of length ℓ .

Suppose that $\varepsilon n/3 < \ell \leq n$. Let C be the properly coloured cycle given by Lemma 4.1, so $|C| \leq 2\varepsilon n/3$. Let G'' be the subgraph of G obtained after removing all the vertices of C . Note that $\delta_1^c(G'') > (2 + \varepsilon)n/3 \geq (2 + \varepsilon)|G''|/3$. By Lemma 3.3, G'' can be covered by at most $6\varepsilon^{-1}$ vertex-disjoint properly coloured paths. That is, we can find vertex-disjoint properly coloured paths $P_1, \dots, P_k$ in G'' such that $k \leq 6\varepsilon^{-1}$, $V(P_i) \cap V(P_j) = \emptyset$ for all $i \neq j$ and $\bigcup P_i = V(G'')$. By removing vertices in the paths, we may assume that the paths $P_1, \dots, P_k$ span exactly $\ell - |C|$ vertices. By the property of C guaranteed by Lemma 4.1, there exists a properly coloured cycle C' with $V(C') = V(C) \cup \bigcup_{1 \leq i \leq k} V(P_i)$. Note that $|C'| = |C| + \sum |P_i| = \ell$. Therefore, C' is a properly coloured cycle of length ℓ . $\square$

6. ACKNOWLEDGEMENTS

The author would like to thank Katherine Staden for her comments on a draft of this paper.

REFERENCES

- [1] N. Alon and G. Gutin, *Properly colored Hamilton cycles in edge-colored complete graphs*, Random Structures Algorithms **11** (1997), 179–186.
- [2] N. Alon and J. H. Spencer, *The probabilistic method*, second ed., Wiley-Interscience Series in Discrete Mathematics and Optimization, Wiley-Interscience [John Wiley & Sons], New York, 2000, With an appendix on the life and work of Paul Erdős.
- [3] J. Bang-Jensen and G. Gutin, *Digraphs*, second ed., Springer Monographs in Mathematics, Springer-Verlag London Ltd., London, 2009, Theory, algorithms and applications.
- [4] J. Bang-Jensen, G. Gutin, and A. Yeo, *Properly coloured Hamiltonian paths in edge-coloured complete graphs*, Discrete Appl. Math. **82** (1998), no. 1-3, 247–250.
- [5] B. Bollobás and P. Erdős, *Alternating Hamiltonian cycles*, Israel J. Math. **23** (1976), 126–131.
- [6] C. C. Chen and D. E. Daykin, *Graphs with Hamiltonian cycles having adjacent lines different colors*, J. Combinatorial Theory Ser. B **21** (1976), no. 2, 135–139.
- [7] J. Feng, H. Giesen, Y. Guo, G. Gutin, T. Jensen, and A. Rafiey, *Characterization of edge-colored complete graphs with properly colored Hamilton paths*, J. Graph Theory **53** (2006), no. 4, 333–346.
- [8] J. W. Grossman and R. Häggkvist, *Alternating cycles in edge-partitioned graphs*, J. Combin. Theory Ser. B **34** (1983), 77–81.
- [9] H. Li and G. Wang, *Color degree and alternating cycles in edge-colored graphs*, Discrete Math. **309** (2009), 4349–4354.
- [10] A. Lo, *A Dirac type condition for properly coloured paths and cycles*, J. Graph Theory, to appear.
- [11] A. Lo, *Properly coloured Hamiltonian cycles in edge-coloured complete graphs*, (submitted).
- [12] V. Rödl, A. Ruciński, and E. Szemerédi, *An approximate Dirac-type theorem for k -uniform hypergraphs*, Combinatorica **28** (2008), no. 2, 229–260.
- [13] J. Shearer, *A property of the colored complete graph*, Discrete Math. **25** (1979), no. 2, 175–178.
- [14] A. Yeo, *A note on alternating cycles in edge-coloured graphs*, J. Combin. Theory Ser. B **69** (1997), 222–225.